
MODULATION OF SODIUM TRANSPORT DURING INTERFERON- α 2b-INDUCED DIFFERENTIATION IN HUMAN NEUROBLASTOMA CELLS

O.V. DOLHA, O.M. ROZHMANOVA, N.Kh. POHORELA, I.S. MAHURA

Bogomoletz Institute of Physiology, Ukraine

dolgaya@biph.kiev.ua



Olena Dolha is a senior researcher of the Department of Neurochemistry at the Bogomoletz Institute of Physiology, Kyiv. She received her Ph.D. in biology in 1977. For the past decade, Dr. Dolga's research has focused on the molecular basis for bidirectional neural-immune interactions. She studies the role of interferon- α in regulation of the nerve cells functions.



Olha Rozhmanova (1935-2007), Ph.D. in biology. She was a senior researcher of the Department for neurochemistry at the Bogomoletz Institute of Physiology, Kyiv. Dr. Rozhmanova had a longstanding and continuing interest in the mechanisms of interferon- α involvement in regulation of ion transport in nerve cells.



Nelly Pohorela is a senior researcher of the Department for general physiology of nerve system at the Bogomoletz Institute of Physiology, Kyiv. She received her Ph.D. in histology and embryology in 1975. Last years, Dr. Pohorela's studies have been devoted to the contemporary cell culturing methods, cell differentiation under various endo- and exogenous factors, and morphological features of the transformed cell lines. She is responsible for the Bank of Cell Lines, which is listed to the State Register of objects constituting National Property of Ukraine.

Information about prof. I.S. Mahura is described on p. 87.

Introduction

Over the past 20 years, neural-immune interactions are a matter of topical interest in modern neurobiological investigations. Recent findings indicate that the central nervous and immune systems are intimately linked. These two systems use the common signals molecules (hormones, neurotransmitters, neuropeptides, cytokines), and common receptors for the bidirectional communication and regulation. In this respect, studies of the neurotropic activity of cytokines released by activated immune cells have been of fundamental importance. According to modern concepts, interferons (IFNs) are one of the key polyfunctional cytokines providing integrative activity of the neural and immune systems (Blalock, 1989; Wrona D., 2006).

IFNs are inducible proteins that regulate resistance to viral infection, enhance innate and acquired immune responses, modulate cellular metabolism, and control cell growth and differentiation. After binding to high-affinity receptor on the cell surface, IFNs initiate a signaling cascade through signaling proteins. Cellular actions are mediated through specific IFN-stimulated genes (ISGs), which underlie a variety of cellular responses (Borden et al., 2007).

IFN- α was originally considered as a cytokine with antiviral, immunomodulating, and antiproliferative properties. Current studies indicate that IFN- α affects the central nervous system activity. It was demonstrated that IFN- α administration results in an alteration of the electrophysiological activity of brain regions participating in pain suppression mechanisms and temperature and food-intake regulation. Some of the reports indicate that IFN- α modulates neuronal activity via opiate receptors (for review, see Dafny; Yang., 2005). It was shown that IFN- α 2b is involved in regulation of neurogenesis (Rozhmanova et al., 2004). These findings suggest that IFN- α is one of the regulatory mediators, which links the central nervous and the immune systems.

Since the disturbance of this interaction underlies many disorders of the central nervous system, the problem of neural-immune interactions is not only of a theoretical importance. Therefore, both experimental and clinical aspects of this interaction are now intensively investigated.

At present, recombinant IFN- α is widely used for treatment of a number of viral and oncological diseases, in particular neuroblastoma (Bruchelt et al., 1990; Gutterman, 1994). The molecular mechanisms of IFN antiproliferative activity are still obscure. The processes of cell growth and division depend significantly on the concentration gradients of Na^+ , K^+ , and Ca^{2+} ions (Vereninov; Marakhova, 1986). Therefore, the antiproliferative effect of IFN- α 2b can be related to the dysfunction of the ion-transporting systems. Taking into account the above arguments, we focused our attention on the changes in transport of sodium ions, which accompany interferon- α 2b-induced differentiation of human neuroblastoma cells.

Morphological differentiation of IMR 32 cells

Human neuroblastoma IMR 32 cell line is a classic model for the investigation of neuronal differentiation induced by different factors, in particular, cytokines. In our experiments we used human recombinant IFN- α 2b (laferon). IMR 32 cells were cultured during twenty-two hours in the modified media containing 600 IU/ml IFN- α 2b. As criteria of morphological differentiation, we used generally accepted parameters for nerve cells, such as a decrease in the proliferative activity of the cells, an increase in the total length of the neuritis, and a change in the projective area of soma.

At the end of the first day of culturing, the differences between control cells and those placed in the modified media were observed (Rozhmanova et al., 2000). Statistical data of changes in cell density demonstrate that IFN- α 2b inhibited cell proliferation and induced differentiation in 50% of the cultured IMR 32 cells. At the same time, the mean value of the projective area of cell soma in differentiated cells was by 56% larger than that of the control cells. Mean values of the total neurite length did not significantly differ from the analogous parameter in the control group. A significant difference in the length of processes was observed 48 h after IFN- α 2b had been applied.

Along with suppression of the proliferative activity and changes in the morphological parameters, the enhancement of protein synthesis in IMR 32 cells was noted. 22 h after IFN- α 2b addition both total protein content in the cell and protein content in the membrane fraction were, on average, 2.3 times those in the control cells. The present results show that IFN- α 2b controls the proliferation and induces differentiation of IMR 32 cells.

Effect of IFN- α 2b on the Na^+ influx

The role of sodium ions transport mechanisms in cell proliferation is still unknown. However, there is increasing evidence that they are involved in regulation of the cell cycle (Arcangeli; Becchetti, 2006). The study of these mechanisms is very important for understanding of certain regulatory pathways, which modulate the transition between proliferative and nonproliferative states. Therefore, we focused our attention on the changes of inward and outward sodium fluxes in IMR 32 cells, which accompany cell differentiation (Rozhmanova et al., 2000, 2001).

Early changes of the Na^+ influx. Using corresponding extensively known techniques, we studied the effects of short-lasting (to 1 h) and long-lasting (22 h) action of IFN α 2b on the $^{22}\text{Na}^+$ influx and activity of the Na^+ , K^+ -ATPase in IMR 32 cells (Rozhmanova, 2001).

Our experimental results convincingly demonstrated that IFN- α 2b modulates $^{22}\text{Na}^+$ influx in IMR 32 cells at the resting potential. It was demonstrated that after 20 min IFN- α 2b application the $^{22}\text{Na}^+$ influx in neuroblastoma cells was reduced by 20%, in comparison with the control cells, and then gradually recovered.

Veratrine increases probability of opening of sodium channels, while the scorpion toxin from *Leiurus quinquestriatus* inhibits their inactivation in neuroblastoma cells (Catterall, 1975). In the presence of these neurotoxins, in 20 min following the onset of IFN- α 2b application, the $^{22}\text{Na}^+$ influx increased by about 50% in the control cells and by 70 % in the cells affected by IFN- α 2b. A blocker of voltage-operated sodium channels, tetrodotoxin (TTX, $4 \cdot 10^{-7}$ M), completely blocked the $^{22}\text{Na}^+$ influx in the control cells and only by about 75% in the IFN- α 2b-treated cells.

These findings allow one to conclude that IFN- α 2b modulates the transport of sodium ions through the voltage-operated TTX-sensitive sodium channels. Our data are in agreement with electrophysiological investigations, which demonstrated that IFN- α 2b alters characteristics of such channels in undifferentiated IMR 32 cells (Kucher et al., 2000).

Delayed changes of the Na^+ influx. In the cells cultured for 22 h in the medium containing 600 IU/ml IFN- α 2b, the $^{22}\text{Na}^+$ influx in the absence of neurotoxins was by 65% larger, and in the presence of neurotoxins it was by 220% larger than that observed in the control (undifferentiated) cells. TTX completely blocked $^{22}\text{Na}^+$ influx through modified with neurotoxins sodium channels in undifferentiated cells, while the $^{22}\text{Na}^+$ influx dropped only by 50% upon the action of TTX in differentiated cells. As these findings indicate the fact that the long-lasting action of IFN- α 2b causes the appearance of voltage-operated TTX-insensitive sodium channels in the plasma membrane of the IMR 32 cells. The role of Na^+ channels in cell differentiation has not been entirely established. However, there are evidences that regulation of the expression and function of sodium channels play important role during cell differentiation (Gomez et al., 1993; Tabb et al., 1994).

Effect of IFN- α 2b on Na^+ , K^+ -ATPase activity

Na^+ , K^+ -ATPase maintains the ion asymmetry in both nonexcitable and excitable cells and is involved in the control of different cell processes, including cell growth, division, and differentiation. An important condition for realization of the entire cycle of cell reproduction is activation of the Na^+ , K^+ -ATPase system, which accompanies transitions of the cell into G_1 phase (Vereninov; Marakhova, 1986). Therefore, we could suggest that IFN- α 2b changes Na^+ , K^+ -ATPase activity during IMR 32 cells differentiation.

The activity of Na^+ , K^+ -ATPase was estimated in a microsomal fraction obtained from undifferentiated (control) or differentiated IMR 32 cells (Rozhmanova et al., 2000). In undifferentiated cells Na^+ , K^+ -ATPase activity decreased by half within 5 min of IFN- α 2b addition. Following 22 h culturing in the IFN- α 2b-containing medium, in differentiated cells, this activity recovered to 80% of its level in undifferentiated cells.

Evaluating the data obtained in our study, we can conclude that IFN- α 2b induced differentiation of human neuroblastoma cells IMR 32. This effect is associ-

ated with modulation of sodium ions transport through voltage-operated sodium channels and Na^+ , K^+ -ATPase activity. The principles learned from *in vitro* studies are very important in our understanding of IFN- α 2b neurotropic actions and can be used to evaluate its antiproliferative effect *in vivo*.

Acknowledgements. We are grateful to Professor P.G. Kostyuk for his support, inspirational guidance, and useful discussions.

REFERENCES

- Arcangeli A, Becchetti A, 2006. Ion channels and the cell cycle. In: Janigro D (Ed.) The cell cycle in the central nervous system, Humana press, pp. 81-94.
- Blalock JE, 1989. A molecular basis for bidirectional communication between the immune and neuroendocrine systems. *Physiol. Rev.* 69: 1-32.
- Borden EC, Sen GC, Uze G, Silverman RH, Ronsohoff RM, Foster GR, Stark GR, 2007. Interferons at age 50: past, current and future impact on biomedicine. *Nature Rev.* 6: 975-990.
- Bruchelt G., Handgretinger R., Shilbach-Struckle K., Shilling FH, Pollwein P, Schwab M, Treuner J, Niethammer D, 1990. The role of interferons in neuroblastoma. I. Antiproliferative effects. *Klin. Padiatr.* 202: 202-205.
- Catterall WA, 1975. Activation of the action potential sodium ionophore of cultured neuroblastoma cells by veratridine and batrachotoxine. *J. Biol. Chem.* 250: 4053-4059.
- Dafny N, Yang PB, 2005. Interferon and the central nervous system. *Eur. J. Pharmacol.* 523: 1-15.
- Gomez MP, Waloga G, Nasi E, 1993. Introduction of voltage-dependent sodium channels by *in vitro* differentiation of human retinoblastoma cells. *J. Neurophysiol.* 70: 1487-1496.
- Guterman JU, 1994. Cytokine therapeutics: lessons from interferon α . *Proc. Natl. Acad. Sci.* 91: 1198-1205.
- Kucher VV, Magura IS, Rozhmanova OM, Dolgaya EV, Pogorelay N, Kh. 2001. Modulation of voltage-gated sodium channels by recombinant interferon IFN- α 2b in human neuroblastoma cells IMR-32. *Ukr. Biochem. J.* 73: 112-115 (in Russian).
- Rozhmanova OM, Dolgaya EV, Kucher VV, Miransky AV, Stelmakh LN, Pogorelay N, Kh., Magura IS, Matzuka GKh, 2000. Sodium transport in the human neuroblastoma cells during early phase of differentiation with recombinant interferon- α 2b (laferon). *Biopolymers Cell.* 16: 540-546 (in Russian).
- Rozhmanova OM, Dolgaya EV, Stelmakh LN, Pogorelay N, Kh., 2001. Effect of recombinant interferon- α 2b (laferon) on transport of sodium ions in human neuroblastoma cells. *Neurophysiology.* 33: 19-22.
- Rozhmanova OM, Dolgaya EV, Stelmakh LN, Vasilovskaya SV, Votyakova IA, Miranskii AV, Magura IS, 2004. Effect of interferon- α 2b on cells from human embryonic nerve tissue in the early stages of neurogenesis. *Neurophysiology.* 36: 319-324.
- Tabb JS, Fanger GR, Wilson EM, Maue RA, Henderson LP, 1994. Suppression of sodium channel function in differentiating C2 muscle cells stably overexpressing rat androgen receptors. *J. Neurosci* 14: 763-773.
- Vereninov AA, Marakhova II, 1986. Ion transport in cultured cells, Leningrad, Nauka, p 292.
- Wrona D, 2006. Neural-immune interactions: An integrative view of the bidirectional relationship between the brain and immune systems. *J. Neuroimmunol.* 172: 38-58.